

Critically-bound four-body molecules

J.-M. Richard

*Institut des Sciences Nucléaires, Université Joseph Fourier – CNRS-IN2P3,
53, avenue des Martyrs, F-38026 Grenoble Cedex, France*

(Dated: February 2, 2008)

The $(p, d, \bar{p}, \bar{d})$ molecule, with a proton, a deuteron and their antiparticles, is stable against spontaneous dissociation, but none of its three-body subsystems are stable. This molecule should be built by combining two atoms, for instance a protonium ($p\bar{p}$) and its heavier analogue ($d\bar{d}$). Most other four-body molecules have at least one stable three-body subsystem and thus can be built by adding the constituents one by one.

Bressanini et al. [1] have studied the stability of four-charge systems with masses (M^+, m^+, M^-, m^-) . For $M = m$, this corresponds to the positronium molecule Ps_2 , whose stability was first demonstrated in 1947 [2]. For $M \gg m$ or $M \ll m$, this is a hydrogen–antihydrogen $\text{H}\bar{\text{H}}$ system (without annihilation, strong interaction, etc.) which hardly competes with the deeply bound protonium (M^+M^-) involved in the lowest threshold $(M^+M^-) + (m^+m^-)$. Stability is thus restricted to an interval of M/m close to unity. The Monte-Carlo calculation of Ref. [1] leads to an estimate

$$\frac{1}{2.2} \lesssim \frac{M}{m} \lesssim 2.2, \quad (1)$$

which is confirmed by a powerful variational method [3].

The case of three unit charges is well documented [4, 5, 6], in particular for the $(M^\pm, m^\mp, m^\pm)$ configurations. For $M = m$, this is the stable positronium ion Ps^- . For $M \gg m$, we have (p, e^-, e^+) , and for $M \ll m$, $(\bar{p}, p, e^-)$, both unstable. Mitroy [7], using the same stochastic variational approach as in Ref. [3], found that stability is confined to

$$0.70 \lesssim M/m \lesssim 1.64. \quad (2)$$

Comparing the results (1) and (2) indicates a window for “Borromean” binding. For instance, for $M/m = 2$, which is the deuteron-to-proton mass ratio, the (M^+, m^+, M^-, m^-) molecule is bound, but neither $(M^\pm, m^\mp, M^\pm)$ nor $(m^\pm, m^\mp, M^\pm)$ are stable.

The word “Borromean” has been proposed in nuclear physics to identify bound states whose subsystems are unbound [8]. It comes from the Borromean rings, which are interlaced in such a subtle topological way, that if any one of them is removed, the two others become unlocked. For instance, the ${}^6\text{He}$ isotope of ordinary Helium is stable, while ${}^5\text{He}$ is not. In a three-body picture, this means that the (α, n, n) system is bound, whereas (α, n) and (n, n) are unbound.

For $N > 3$ constituents, one might define Borromean binding as the property of all N' -body subsystems being unstable, with $N' = 2$, or $N' = N - 1$, or $N' < N$. We propose the following definition: A bound state is *Borromean* if there is no path to build the system via a series of stable bound states by adding the constituents one by one. Then, $(p, d, \bar{p}, \bar{d})$ is Borromean. It is truly an atom–atom composite, more representative of larger molecules of ordinary chemistry. The same is true for neighboring systems $(m_1^+, m_2^+, m_3^-, m_4^-)$ with less symmetry. A minimal extension of the domain of stability can be derived using the variational principle [9].

In comparison, H_2 or Ps_2 systems appear to be more robust, with several three-body subsystems being stable, (p, e^+, e^-) or (p, p, e^-) for H_2 , and $(e^\pm, e^\mp, e^\mp)$ for Ps_2 . The positronium hydride PsH (p, e^+, e^-, e^-) contains the unstable (p, e^+, e^-) , but also the stable (p, e^-, e^-) and (e^+, e^-, e^-) , and thus is not Borromean.

Note that if the antideuteron is replaced by the celebrated Ω^- hyperon (predicted by Gell-Mann by symmetry considerations which led to the quark model, and discovered by Samios et al. at Brookhaven [10]), and if the deuteron is replaced by $\bar{\Omega}^+$, the mass ratio $M/m = 1.78$ becomes close to one of the critical values of Eq. (2). If $A = (\Omega^- \bar{\Omega}^+)$, we have an effective $(A, p, \bar{p})$ three-body system with both (A, p) and $(A, \bar{p})$ energies vanishing. The Efimov effect [11] survives finite-size effects, since it is governed by the long range part of the interaction. However, the Coulomb attraction between p and $\bar{p}$ spoils the $-1/\rho^2$ behavior (ρ is the hyperradius) necessary in the hypercentral potential for Efimov states to appear. See, e.g., the approach by Fedorov and Jensen, in Ref. [11].

This investigation grew out of discussions with Dario Bressanini and André Martin. Comments by E.A.G. Armour, A.J. Cole, A.S. Jensen and K. Varga are also gratefully acknowledged. A partial and preliminary version of this comment was presented at the Few-Body Conference in Bled [9].

[1] D. Bressanini, M. Mella and G. Morosi, Phys. Rev. A **55**, 200 (1997).

[2] E.A. Hylleraas and A. Ore, Phys. Rev. **71**, 493 (1947).

[3] K. Varga, Few-Body Systems Suppl. **10**, 11 (1999) (

- Proc. 16th European Conference on Few-Body Problems in Physics, Autrans, France, June 1998, ed. B. Desplanques et al.).
- [4] A. Martin, J.-M. Richard and T.T. Wu, Phys. Rev. **A43**, 3648 (1992).
 - [5] E.A.G. Armour and W. Byers Brown, Accounts of Chemical Research, **26**, 168 (1993).
 - [6] E.A.G. Armour, J.-M. Richard, and K. Varga, “Stability of Few-Charge Systems”, in preparation.
 - [7] J. Mitroy, J. Phys. B: At. Mol. Opt. Phys. **33** 5307 (2000).
 - [8] M.V. Zhukov, B.V. Danilin, D.V. Fedorov, J.M. Bang, I.S. Thompson, and J.S. Vaagen, Phys. Rep. **231** 151 (1993).
 - [9] J.-M. Richard, Talk at the XVIII European Few-Body Conference, Bled, Slovenia, Sept. 2002, to appear in Few-Body Systems Suppl., ed. M. Rosina et al.
 - [10] See, for instance, A. Pais, *Inward Bound, of Matter and Forces in the Physical World* (Clarendon Press, Oxford, 1986).
 - [11] V. Efimov, Phys. Lett. **33B** (1970) 560; Sov. J. Nucl. Phys. **12** (1971) 581; R.N. Amado and R.V. Noble, Phys. Lett. **35B** (1971) 25; Phys. Rev. D **5** (1972) 1992; S. Albeverio, R. Høegh-Krohn and T.T. Wu, Phys. Lett. **83A** (1981) 101; D.V. Fedorov and A.S. Jensen, Phys. Rev. Lett. **71** (1993) 4103.